

SOLUTION OF VARIATIONAL INEQUALITY PROBLEMS ON FIXED POINT SETS OF NONEXPANSIVE MAPPINGS USING ITERATIVE METHODS.

ERIC U. OFOEDU*,¹

ABSTRACT. In this paper, we introduce new implicit and explicit iterative schemes which converge strongly to a unique solution of variational inequality problems for strongly accretive operators over a common fixed point set of finite family of nonexpansive mappings in q -uniformly smooth real Banach spaces. As an application, we introduce an iteration process which converges strongly to a solution of the variational inequality which is a common fixed point of finite family of strictly pseudocontractive mappings. Our theorems extend, generalize, improve and unify the corresponding results of Xu [48] and Yamada [50] and that of a host of other authors. Our corollaries and our method of proof are of independent interest.

1. INTRODUCTION

Let $(E, \|\cdot\|)$ be a real normed space. Let $S := \{x \in E : \|x\| = 1\}$. The space E is said to have a *Gâteaux differentiable* norm if the limit

$$\lim_{t \rightarrow 0} \frac{\|x + ty\| - \|x\|}{t}$$

exists for each $x, y \in S$. E is said to have a *uniformly Gâteaux differentiable* norm if for each $y \in S$ the limit is attained uniformly for $x \in S$. The space E is said to be *uniformly smooth* if and only if for all $\epsilon > 0$, there exists $\delta > 0$ such that for all $x, y \in E$ with $\|x\| = 1$ and $\|y\| \leq \delta$, the inequality

$$\frac{\|x + y\| + \|x - y\|}{2} - 1 < \epsilon\|y\|$$

holds. It is well known that every uniformly smooth real Banach space is a *reflexive real Banach space* and has uniformly Gâteaux differentiable norm (see e.g., [13]).

Let E be a real normed linear space with dimension, $\dim(E) \geq 2$. The *modulus of smoothness* of E is the function $\rho_E : [0, \infty) \rightarrow [0, \infty)$ defined by

$$\rho_E(t) = \sup \left\{ \frac{\|x + y\| + \|x - y\|}{2} - 1 : \|x\| = 1, \|y\| = t \right\}.$$

Key words and phrases. Modulus of smoothness, generalized duality maps, accretive operators, strongly accretive mappings, Strictly pseudocontractive mappings, uniformly Gâteaux differentiable norm.

2000 *Mathematics Subject Classification.* 47H06, 47H09, 47J05, 47J25.

*The author undertook this work when he visited the Abdus Salam International Centre for Theoretical Physics (ICTP), Trieste, Italy as a visiting fellow.

In terms of the modulus of smoothness (see e.g. [13]), the space E is called *uniformly smooth* if and only if $\lim_{t \rightarrow 0^+} \frac{\rho_E(t)}{t} = 0$. E is called *q -uniformly smooth* if and only if there exists a constant $c > 0$ such that $\rho_E(t) \leq ct^q$, $t > 0$. It is easy to see that for $1 < q < +\infty$, every q -uniformly smooth Banach space is uniformly smooth; and thus has a uniformly Gâteaux differentiable norm.

Let E be a real normed linear space with dual E^* . We denote by J_q the *generalized duality mapping* from E to 2^{E^*} defined by

$$J_q x := \{f^* \in E^* : \langle x, f^* \rangle = \|x\|^q, \|f^*\| = \|x\|^{q-1}\},$$

where $\langle \cdot, \cdot \rangle$ denotes the generalized duality pairing between members of E and members of E^* . For $q = 2$, the mapping $J = J_2$ from E to 2^{E^*} is called *the normalized duality mapping*. It is well known that if E is uniformly smooth or E^* is strictly convex, then duality mapping is single-valued; and if E has a uniformly Gâteaux differentiable norm then the duality mapping is *norm-to-weak* uniformly continuous* on bounded subsets of E . If $E = H$ is a Hilbert space then the duality mapping becomes the identity map of H (see e.g., [13, 52]). In the sequel, we shall denote the single-valued generalized duality mapping by j_q and the single valued normalized duality mapping by j .

A real normed space E with strictly convex dual is said to have a *weakly sequentially continuous generalized duality mapping* j_q if and only if for each sequence $\{x_n\}_{n \geq 1}$ in E such that $\{x_n\}_{n \geq 1}$ converges weakly to x^* in E , we have that $\{j_q(x_n)\}_{n \geq 1}$ converges in the weak* topology to $j_q(x^*)$. A real Banach space E is said to satisfy *Opial's condition* if for any sequence $\{x_n\}_{n \geq 1}$ in E such that $\{x_n\}_{n \geq 1}$ converges weakly to x^* in E , we have that

$$\limsup_{n \rightarrow \infty} \|x_n - x^*\| < \limsup_{n \rightarrow \infty} \|x_n - y\|$$

for all $y \in E$, $y \neq x^*$. By Theorem 1 of Gossez and Lami Dozo [17], it is known that if E admits weakly sequentially continuous duality mapping, then E satisfies Opial's condition (see [17] for more details). Aside from Hilbert spaces, it was noted in [5] that the most significant class of Banach spaces having a weakly sequentially continuous generalized duality mapping are the sequence spaces ℓ_q for $1 < q < +\infty$ (see [6] and [7], where it is also shown that $L_q(\mathbb{R})$ has no weakly sequentially continuous duality mapping for $q \neq 2$).

A mapping $f : E \rightarrow E$ is said to be a *strict contraction* or simply a *contraction* if and only if there exists $\gamma_0 \in [0, 1)$ such that for all $x, y \in E$,

$$\|f(x) - f(y)\| \leq \gamma_0 \|x - y\|.$$

A mapping $T : E \rightarrow E$ is called *nonexpansive* if and only if for all $x, y \in E$,

$$\|Tx - Ty\| \leq \|x - y\|.$$

A point $x \in E$ is called a *fixed point* of an operator T if and only if $Tx = x$. The set of fixed points of an operator T is denoted by $F(T)$, that is, $F(T) := \{x \in E : Tx = x\}$.

A mapping $A : E \rightarrow E$ is said to be *accretive* if and if for all $x, y \in E$ there exists $j_q(x - y) \in J_q(x - y)$ such that

$$\langle Ax - Ay, j_q(x - y) \rangle \geq 0.$$

The operator $A : E \rightarrow E$ is said to be *ψ -uniformly accretive* if there exists a strictly increasing continuous function $\psi : [0, +\infty) \rightarrow [0, +\infty)$ with $\psi(0) = 0$ such that for all $x, y \in E$, there exists $j_q(x - y) \in J_q(x - y)$ such that

$$\langle Ax - Ay, j_q(x - y) \rangle \geq \psi(\|x - y\|).$$

$A : E \rightarrow E$ is called *strongly accretive* if and only if there exists a constant $\eta > 0$ and for all $x, y \in E$ there exist $j_q(x - y) \in J_q(x - y)$ such that

$$\langle Ax - Ay, j_q(x - y) \rangle \geq \eta \|x - y\|^q.$$

It is easy to see that every strongly accretive mapping is ψ -uniformly accretive with $\psi(\tau) = \eta\tau^q \forall \tau \in [0, +\infty)$. When $E = H$ is a Hilbert space, accretive, ψ -uniformly accretive, strongly accretive mappings coincide with *monotone*, *ψ -uniformly monotone*, *strongly monotone* mappings, respectively.

A bounded linear operator $A : H \rightarrow H$ is called a *strongly positive* operator if and only if there exist a constant $k > 0$ such that for all $x \in H$,

$$\langle Ax, x \rangle \geq k \|x\|^2.$$

Thus, every strongly positive bounded linear operator on H is strongly monotone.

Iterative approximation of fixed points and zeros of nonlinear operators has been studied extensively by many authors to solve nonlinear operator equations as well as variational inequality problems (see e.g., [20], [25]-[31], [35]-[41], [14, 15]). In particular, iterative approximation of fixed points of nonexpansive mappings is an important subject in nonlinear operator theory and its applications in image recovery and signal processing is well known (see e.g., [12, 34, 51]). Most published results on nonexpansive mappings centered on the iterative approximation of fixed points of the nonexpansive mappings or approximation of a common fixed points of a given family of nonexpansive mappings.

Iterative methods for nonexpansive mappings are now also applicable in solving convex minimization problems (see, for example, [48] and references therein). Let H be a real Hilbert space with inner product $\langle \cdot, \cdot \rangle$. Let C be a closed convex nonempty subset of H , let $T : C \rightarrow C$ be a nonexpansive mapping such that $F(T) \neq \emptyset$. Given $u \in C$ and a real sequence $\{\alpha_n\}_{n \geq 1}$ in the interval $(0, 1)$, starting with an arbitrary initial $x_0 \in C$, let a sequence $\{x_n\}_{n \geq 1}$ be defined by

$$(1.1) \quad x_{n+1} = \alpha_{n+1}u + (1 - \alpha_{n+1})Tx_n, n \geq 0.$$

Under appropriate conditions on the iterative parameter $\{\alpha_n\}_{n \geq 1}$, it has been shown by Halpern [18], Lions [23], Wittmann [44] and Bauschke [2] that $\{x_n\}_{n \geq 0}$ converges strongly to $P_{F(T)}u$, the projection of u to the fixed point set, $F(T)$ of T . This means that the limit of the sequence $\{x_n\}_{n \geq 0}$ solves the following minimization

problem:

$$\text{find } x^* \in F(T) \text{ such that } \|x^* - u\| = \min_{x \in F(T)} \|x - u\|.$$

H. K. Xu [48] studied the following quadratic minimization problem: find $x^* \in F(T)$ such that

$$(1.2) \quad \frac{1}{2} \langle Ax^*, x^* \rangle - \langle x^*, u \rangle = \min_{x \in F(T)} \left(\frac{1}{2} \langle Ax, x \rangle - \langle x, u \rangle \right),$$

where $u \in H$ is fixed and $A : H \rightarrow H$ a bounded linear strongly positive operator. Let $C_1, C_2, \dots, C_N$ be N closed convex subsets of a real Hilbert space H having a nonempty intersection C . Suppose also that each C_i is a fixed point set of nonexpansive mappings $T_i : H \rightarrow H, i = 1, 2, \dots, N$, Xu [48] proved strong convergence of the iterative algorithm

$$(1.3) \quad x_0 \in H, x_{n+1} = (I - \alpha_{n+1}A)T_{n+1}x_n + \alpha_{n+1}u, n \geq 0$$

(where $T_n = T_{n \bmod N}$ and the mod function takes values in $\{1, 2, \dots, N\}$) to a unique solution of the quadratic minimization problem (1.2).

Marino and Xu [26], proved that the iteration scheme given by

$$(1.4) \quad x_0 \in H, x_{n+1} = \alpha_n \gamma f(x_n) + (I - \alpha_n A)T x_n, n \geq 0$$

converges strongly to a unique solution $x' \in F(T)$ of the variational inequality

$$(1.5) \quad \langle (\gamma f - A)x', y - x' \rangle \leq 0 \quad \forall y \in F(T),$$

which is the optimality condition for the minimization problem

$$\min_{x \in F(T)} \left(\frac{1}{2} \langle Ax, x \rangle - h(x) \right),$$

where h is a potential function for γf (that is, $h'(x) = \gamma f(x)$ for all $x \in H$); provided $f : H \rightarrow H$ is a contraction, $T : H \rightarrow H$ is nonexpansive and the iterative parameter $\{\alpha_n\}_{n \geq 0}$ satisfies appropriate conditions.

In [50], Yamada introduced the following hybrid iterative method

$$(1.6) \quad x_0 \in H, x_{n+1} = T x_n - \mu \lambda_n A(T x_n), n \geq 0,$$

where T is nonexpansive, A is L -Lipschitzian and strongly monotone operator with constant $\eta > 0$ and $0 < \mu < \frac{2\eta}{L^2}$. He proved that if $\{\lambda_n\}_{n \geq 1}$ satisfy appropriate conditions, then (1.6) converges strongly to a unique solution $x' \in F(T)$ of the variational inequality

$$\langle Ax', y - x' \rangle \geq 0 \quad \forall y \in F(T).$$

Recently, M. Tian [43] introduced the following iterative method:

$$(1.7) \quad x_0 \in H, x_{n+1} = \alpha_n \gamma f(x_n) + (I - \mu \alpha_n A)T x_n, n \geq 0.$$

Tian [43] proved that if $f : H \rightarrow H$ is a contraction, $A : H \rightarrow H$ is an η -strongly monotone mapping, $T : H \rightarrow H$ a nonexpansive mapping and the parameter $\{\alpha_n\}_{n \geq 1}$ satisfies appropriate conditions, then the sequence $\{x_n\}_{n \geq 1}$ converges strongly to a unique solution $x' \in F(T)$ of the variational inequality

$$\langle (\gamma f - \mu A)x', y - x' \rangle \leq 0 \quad \forall y \in F(T).$$

In this paper, motivated by the results of the authors mentioned above, it is our aim to introduce new implicit and explicit iterative schemes which converge strongly to a unique solution of variational inequality problem over a common fixed point set of finite family of nonexpansive mappings in q -uniformly smooth real Banach spaces. As an application, we introduce an iteration process which converges strongly to a solution of the variational inequality which is a common fixed point of family of strictly pseudocontractive mappings. Our theorems extend, generalize, improve and unify the corresponding results of Xu [48] and Yamada [50] and that of a host of other authors. Our corollaries and our method of proof are of independent interest.

2. PRELIMINARY

Let μ be a bounded linear functional defined on ℓ_∞ satisfying $\|\mu\| = 1 = \mu(1)$. It is known that μ is a mean on $\mathbb{N}$ if and only if

$$\inf\{a_n : n \in \mathbb{N}\} \leq \mu(a_n) \leq \sup\{a_n : n \in \mathbb{N}\}$$

for every $a = (a_1, a_2, a_3, \dots) \in \ell_\infty$. In the sequel, we shall use $\mu_n(a_n)$ instead of $\mu(a)$. A mean μ on $\mathbb{N}$ is called a *Banach limit* if $\mu_n(a_n) = \mu_n(a_{n+1})$ for every $a = (a_1, a_2, a_3, \dots) \in \ell_\infty$. It is well known that if μ is a Banach limit, then

$$\liminf_{n \rightarrow \infty} a_n \leq \mu_n a_n \leq \limsup_{n \rightarrow \infty} a_n$$

for all $a = (a_1, a_2, a_3, \dots) \in \ell_\infty$. Furthermore, if $a = (a_1, a_2, \dots)$, $b = (b_1, b_2, \dots) \in \ell_\infty$ and $\lim_{n \rightarrow \infty} a_n = a^*$ ($\lim_{n \rightarrow \infty} (a_n - b_n) = 0$), then $\mu_n(a_n) = a^*$ (respectively, $\mu_n(a_n) = \mu_n(b_n)$).

In what follows, we shall make use of the following Lemmas:

Lemma 2.1. *Let E be a real normed linear space, then for $1 < q < +\infty$, the following inequality holds:*

$$\|x + y\|^q \leq \|x\|^q + q \langle y, j_q(x + y) \rangle \quad \forall x, y \in E, \quad \forall j_q(x + y) \in J_q(x + y).$$

Lemma 2.2. *(See e.g., [4, 46, 48]) Let $\{\lambda_n\}_{n \geq 1}$ be a sequence of nonnegative real numbers satisfying the condition*

$$\lambda_{n+1} \leq (1 - \alpha_n)\lambda_n + \sigma_n, \quad n \geq 0,$$

where $\{\alpha_n\}_{n \geq 0}$ and $\{\sigma_n\}_{n \geq 0}$ are sequences of real numbers such that $\{\alpha_n\}_{n \geq 1} \subset [0, 1]$, $\sum_{n=1}^{\infty} \alpha_n = +\infty$. Suppose that $\sigma_n = o(\alpha_n)$, $n \geq 0$ (i.e., $\lim_{n \rightarrow \infty} \frac{\sigma_n}{\alpha_n} = 0$) or $\sum_{n=1}^{\infty} |\sigma_n| < +\infty$ or $\limsup_{n \rightarrow \infty} \frac{\sigma_n}{\alpha_n} \leq 0$, then $\lambda_n \rightarrow 0$ as $n \rightarrow \infty$.

Lemma 2.3. *(Compare with Lemma 3 pg. 257 of Bruck [11]) Let C be a nonempty closed and convex subset of a real strictly convex Banach space E . Let $\{T_i\}_{i \geq 1}$ be a sequence of nonself nonexpansive mappings $T_i : C \rightarrow E$ such that $F := \bigcap_{i=1}^{\infty} F(T_i) \neq$*

$\emptyset$. Let $\{\sigma_i\} \subset (0, 1)$ be such that $\sum_{i=1}^{\infty} \sigma_i = 1$. Then the mapping $T := \sum_{i=1}^{\infty} \sigma_i T_i : C \rightarrow$

E is well defined, nonexpansive and $F(T) = \bigcap_{i=1}^{\infty} F(T_i)$.

Lemma 2.4. (See [53], p.202 Lemma 3). Let E be a strictly convex Banach space and C be a closed convex subset of E . Let $T_1, T_2, \dots, T_r$ be nonexpansive mappings of C into itself such that the set of common fixed points of $T_1, T_2, \dots, T_r$ is nonempty. Let $S_1, S_2, \dots, S_r$ be mappings of C into itself given by $S_i = (1 - \gamma_i)I + \gamma_i T_i$ for any $0 < \gamma_i < 1$, $i = 1, 2, \dots, r$, where I denotes the identity mapping on C . Then $S_1, S_2, \dots, S_r$ satisfies the following: $\bigcap_{i=1}^r F(S_i) = \bigcap_{i=1}^r F(T_i)$ and $\bigcap_{i=1}^r F(S_i) = F(S_r S_{r-1} \dots S_1) = F(S_1 S_{r-1} \dots S_2) = \dots = F(S_{r-1} \dots S_1 S_r)$.

Lemma 2.5. (See e.g. [47]) Let E be a q -uniformly smooth real Banach space for some $q > 1$, then there exists some positive constant d_q such that

$$(2.1) \quad \|x + y\|^q \leq \|x\|^q + q \langle y, j_q(x) \rangle + d_q \|y\|^q \quad \forall x, y \in E, \quad \forall j_q(x) \in J_q(x).$$

If E is L_q (or ℓ_q) space, the constant d_q in (2.1) has been calculated. This is shown in the following lemma.

Lemma 2.6. (See e.g. [21]) Let E be L_q (or ℓ_q) space ($1 < q < +\infty$) and $x, y \in E$,

(1) if $1 < q < 2$, then

$$\|x + y\|^q \leq \|x\|^q + q \langle y, j_q(x) \rangle + d_q \|y\|^q \quad \forall x, y \in E, \quad \forall j_q(x) \in J_q(x),$$

where $d_q = \frac{1+b_q^{q-1}}{(1+b_q)^{q-1}}$, and b_q is the unique solution of the equation $(q-2)b_q^{q-1} + (q-1)b_q^{q-2} - 1 = 0$, $0 < b < 1$.

(2) if $2 \leq q < +\infty$, then

$$\|x + y\|^2 \leq \|x\|^2 + 2 \langle y, j(x) \rangle + (q-1) \|y\|^2 \quad \forall x, y \in E, \quad \forall j(x) \in J(x).$$

Lemma 2.7. (Lemma 2.2 of [45], p. 1411) Let C be a closed convex nonempty subset of a reflexive Banach space which satisfies Opial's condition and suppose $T : C \rightarrow E$ is nonexpansive, then the mapping $I - T$ is demiclosed at zero, that is $\{x_n\}_{n \geq 1}$ is a sequence in C such that $x_n \rightarrow x^*$ and $x_n - Tx_n \rightarrow 0$ as $n \rightarrow \infty$, then $x^* = Tx^*$.

3. MAIN RESULTS.

Lemma 3.1. Let E be a real normed space. Let $A : E \rightarrow E$ be a strongly accretive mapping with a constant $\eta > 0$. Let $\lambda > 0$, then the mapping $\lambda A : E \rightarrow E$ is strongly accretive.

proof. Observe that

$$(3.1) \quad \begin{aligned} \langle \lambda Ax - \lambda Ay, j_q(x - y) \rangle &= \lambda \langle Ax - Ay, j_q(x - y) \rangle \\ &\geq \lambda \eta \|x - y\|^q. \end{aligned}$$

This completes the proof. $\square$

Lemma 3.2. *Let E be a real normed space. Let $A : E \rightarrow E$ be a strongly accretive mapping with a constant $\eta > 0$. Let $T : E \rightarrow E$ be a nonexpansive mapping such that $F(T) \neq \emptyset$, $\lambda > 0$ and $u \in E$ be a fixed vector. Suppose that the solution $x' \in F(T)$ of the variational inequality $\langle u - \lambda Ax', j_q(p - x') \rangle \leq 0 \ \forall p \in F(T)$ exists, then x' is unique.*

Proof. Suppose for contradiction that the variational inequality has two solutions in $F(T)$, say $x' \neq y'$, then we have, in particular, that

$$\langle u - \lambda Ax', j_q(y' - x') \rangle \leq 0$$

and

$$-\langle u - \lambda Ay', j_q(y' - x') \rangle \leq 0.$$

Thus, adding these two inequalities and using (3.1), we obtain that

$$\lambda \eta \|y' - x'\|^q \leq \langle \lambda Ay' - \lambda Ax', j_q(y' - x') \rangle \leq 0,$$

a contradiction. Hence, the variational inequality has a unique solution, provided the solution exists. This completes the proof. $\square$

Let $1 < q < +\infty$ and let E be a q -uniformly smooth real Banach space. Let $A : E \rightarrow E$ be an L -Lipschitzian strongly accretive mapping with a constant $\eta > 0$. Let $T : E \rightarrow E$ be a nonexpansive mapping. Then for the map $(I - t\lambda A)T : E \rightarrow E$ (where I is the identity map of E , $\lambda > 0$ and $t \in (0, 1)$), we obtain using Lemma 2.5 that

$$\begin{aligned} \|(I - t\lambda A)Tx - (I - t\lambda A)Ty\|^q &= \|Tx - Ty - t\lambda(ATx - ATy)\|^q \\ &\leq \|Tx - Ty\|^q - qt\lambda \langle ATx - ATy, j_q(Tx - Ty) \rangle \\ &\quad + d_q(t\lambda)^q \|ATx - ATy\|^q \\ &\leq \|Tx - Ty\|^q - qt\eta\lambda \|Tx - Ty\|^q \\ &\quad + d_q(tL\lambda)^q \|Tx - Ty\|^q \\ &\leq \left(1 - t\lambda(q\eta - d_q L^q \lambda^{q-1})\right) \|Tx - Ty\|^q \end{aligned}$$

So that if λ is such that $0 < \lambda < \left(\frac{q\eta}{L^q d_q}\right)^{\frac{1}{q-1}}$, we have that

$$0 < 1 - t\lambda(q\eta - d_q L^q \lambda^{q-1}) < 1$$

and since T is a nonexpansive mapping of E into E and $1 < q < +\infty$, we obtain

$$\begin{aligned} \|(I - t\lambda A)Tx - (I - t\lambda A)Ty\| &\leq \left(1 - t\lambda(q\eta - d_q L^q \lambda^{q-1})\right)^{\frac{1}{q}} \|Tx - Ty\| \\ (3.2) \qquad \qquad \qquad &\leq \left(1 - t\lambda(q\eta - d_q L^q \lambda^{q-1})\right)^{\frac{1}{q}} \|x - y\|. \end{aligned}$$

Lemma 3.3. *Let E be a q -uniformly smooth real Banach space. Let $A : E \rightarrow E$ be an L -Lipschitzian strongly accretive mapping with a constant $\eta > 0$. Let $T : E \rightarrow E$ be a nonexpansive mapping such that $F(T) \neq \emptyset$. Let $u \in E$ be fixed,*

$0 < \lambda < \left(\frac{q\eta}{L^q d_q}\right)^{\frac{1}{q-1}}$ and $t \in \left(0, \min\left\{1, \frac{1}{q\eta\lambda - d_q(L\lambda)^q}\right\}\right)$, then there exists a unique $z_t \in E$ such that

$$(3.3) \quad z_t = tu + (I - t\lambda A)Tz_t.$$

Proof. For each $t \in \left(0, \min\left\{1, \frac{1}{q\eta\lambda - d_q(L\lambda)^q}\right\}\right)$, define $S_t : E \rightarrow E$ by $S_tx = tu + (I - t\lambda A)Tx$ for all $x \in E$. Then,

$$\begin{aligned} \|S_tx - S_ty\| &= \|(I - t\lambda A)Tx - (I - t\lambda A)Ty\| \\ &\leq \left(1 - t\lambda(q\eta - d_q(L)^q\lambda^{q-1})\right)^{\frac{1}{q}} \|x - y\| \end{aligned}$$

Thus, for all $t \in \left(0, \min\left\{1, \frac{1}{q\eta\lambda - d_q(L\lambda)^q}\right\}\right)$, we have that S_t is a strict contraction on E . Hence, there exists a unique $z_t \in E$ which satisfies (3.3). This completes the proof. $\square$

Lemma 3.4. Let E , A , $u \in E$ and T be as in Lemma 3.3. Let $\{z_t\}$ satisfy (3.3), then

- (1) The mapping from $\left(0, \min\left\{1, \frac{1}{q\eta\lambda - d_q(L\lambda)^q}\right\}\right)$ to E given by $t \mapsto z_t$ is continuous.
- (2) $\{z_t\}$ is bounded;
- (3) $\lim_{t \rightarrow 0} \|z_t - Tz_t\| = 0$;

Proof.

- (1) Let $t_0 \in \left(0, \min\left\{1, \frac{1}{q\eta\lambda - d_q(L\lambda)^q}\right\}\right)$ be arbitrary. It is enough to show that the mapping from $\left(0, \min\left\{1, \frac{1}{q\eta\lambda - d_q(L\lambda)^q}\right\}\right)$ to E given by $t \mapsto z_t$ is continuous at t_0 . Now, for some constant $M_1 > 0$ and using Lemma 2.1, we have that

$$\begin{aligned} \|z_t - z_{t_0}\|^q &= \|(t - t_0)u - (t - t_0)\lambda ATz_t + (I - t_0\lambda A)Tz_t - (I - t_0\lambda A)Tz_{t_0}\|^q \\ &\leq \|(I - t_0\lambda A)Tz_t - (I - t_0\lambda A)Tz_{t_0}\|^q \\ &\quad + q \left\langle (t - t_0)u - (t - t_0)\lambda ATz_t, j_q(z_t - z_{t_0}) \right\rangle \\ (3.4) \quad &\leq \left(1 - t_0(q\eta\lambda - d_q(L\lambda)^q)\right) \|z_t - z_{t_0}\|^q + |t - t_0| M_1 \|z_t - z_{t_0}\|^{q-1}. \end{aligned}$$

Thus, we obtain from (3.4) that

$$\|z_t - z_{t_0}\| \leq \frac{M_1}{t_0(q\eta\lambda - d_q(L\lambda)^q)} |t - t_0|$$

and the result follows.

- (2) Let $p \in F(T)$. If $z_t = p$ for all $t \in \left(0, \min\left\{1, \frac{1}{q\eta\lambda - d_q(L\lambda)^q}\right\}\right)$, then we are done. Otherwise, let $t^* \in \left(0, \min\left\{1, \frac{1}{q\eta\lambda - d_q(L\lambda)^q}\right\}\right)$ be such that $z_t \neq p \forall t \in (0, t^*]$; then using (3.2), (3.3) and Lemma 2.1, we have that $\forall t \in$

$$\begin{aligned}
 (0, t^*], \\
 \|z_t - p\|^q &= \|(I - t\lambda A)Tz_t - (I - \lambda A)p + t(u - \lambda Ap)\|^q \\
 &\leq \left(1 - t(q\eta\lambda - d_q(L\lambda)^q)\right) \|z_t - p\|^q \\
 &\quad + qt \langle u - \lambda Ap, j_q(z_t - p) \rangle \\
 &\leq \left(1 - t(q\eta\lambda - d_q(L\lambda)^q)\right) \|z_t - p\|^q \\
 &\quad + qt \|u - \lambda Ap\| \cdot \|z_t - p\|^{q-1}
 \end{aligned}
 \tag{3.5}$$

Thus, we obtain from (3.5) that

$$\|z_t - p\| \leq \left(1 - t(q\eta\lambda - d_q(L\lambda)^q)\right) \|z_t - p\| + qt \|u - \lambda Ap\|.
 \tag{3.6}$$

Inequality (3.6) therefore gives

$$\|z_t - p\| \leq \frac{qt \|u - \lambda Ap\|}{q\eta\lambda - d_q(L\lambda)^q}.$$

Hence, the path $\{z_t\}$ is bounded; and so is $\{ATz_t\}$.

(3) Therefore, using (3.3), we have that for some constant $M_0 > 0$,

$$\|z_t - Tz_t\| \leq t \|u - \lambda ATz_t\| \leq tM_0 \rightarrow 0 \text{ as } t \rightarrow 0.$$

This completes the proof. $\square$

Theorem 3.5. *Let E , A , $u \in E$ and T be as in Lemma 3.3. Let $\{z_t\}$ satisfy (3.3), then $\{z_t\}$ converges strongly to some $x' \in F(T)$ which is a solution of the variational inequality*

$$\langle u - \lambda Ax', j(p - x') \rangle \leq 0 \quad \forall p \in F(T).
 \tag{3.7}$$

Proof. By Lemma 3.2, if solution of (3.7) exists in $F(T)$, then it is unique. Let $p \in F(T)$, then using (3.3), we have that

$$z_t - p = t(u - \lambda Ap) + (I - t\lambda A)Tz_t - (I - t\lambda A)p.
 \tag{3.8}$$

Thus, using (3.2), (3.8) and Lemma 2.1,

$$\begin{aligned}
 \|z_t - p\|^q &= \|(I - t\lambda A)Tz_t - (I - t\lambda A)p\|^q \\
 &\quad + qt \langle u - \lambda Ap, j_q(z_t - p) \rangle \\
 &\leq \left(1 - t(q\eta\lambda - d_q(\lambda L)^q)\right) \|z_t - p\|^q \\
 &\quad + qt \langle u - \lambda Ap, j_q(z_t - p) \rangle.
 \end{aligned}
 \tag{3.9}$$

So, (3.9) implies that

$$\|z_t - p\|^q \leq \frac{1}{q\eta\lambda - d_q(\lambda L)^q} \langle u - \lambda Ap, j_q(z_t - p) \rangle.
 \tag{3.10}$$

From Lemma 3.4 we know that $\{z_t\}$ is bounded. Let $\{t_n\}_{n \geq 1}$ in $\left(0, \min\left\{1, \frac{1}{q\eta\lambda - d_q(\lambda L)^q}\right\}\right)$ be such that $\lim_{n \rightarrow \infty} t_n = 0$ and set $z_n = z_{t_n}$. Then, defining $\Psi : E \rightarrow \mathbb{R} \cup \{+\infty\}$ by

$\Psi(x) = \mu_n \|z_n - x\|^q$ for all $x \in E$, where μ_n is a Banach limit on ℓ_∞ , we have that Ψ is continuous, convex and $\lim_{\|x\| \rightarrow +\infty} \Psi(x) = +\infty$. Thus, setting

$$K = \left\{ y \in E : \Psi(y) = \min_{x \in E} \Psi(x) \right\},$$

then K is bounded closed convex and nonempty subset of E and since by Lemma 3.4, $\lim_{n \rightarrow \infty} \|z_n - Tz_n\| = 0$, we obtain using Lemma 2.1 that for some constant $Q^* > 0$ and for all $y \in K$,

$$\begin{aligned} \Psi(Ty) &= \mu_n \|z_n - Ty\|^q = \mu_n \|Tz_n - Ty + z_n - Tz_n\|^q \\ &\leq \mu_n \left(\|Tz_n - Ty\|^q + q \left\langle z_n - Tz_n, j_q(z_n - Ty) \right\rangle \right) \\ &\leq \mu_n \left(\|Tz_n - Ty\|^q + \|z_n - Tz_n\| Q^* \right) \\ (3.11) \quad &\leq \mu_n \|z_n - y\|^q + \mu_n \|z_n - Tz_n\| Q^* = \mu_n \|z_n - y\|^q = \Psi(y). \end{aligned}$$

Thus, $T(K) \subset K$; that is, K is invariant under T . Since E is uniformly smooth real Banach space and thus every bounded closed convex nonempty subset of E has the fixed point property for nonexpansive mappings, then there exists $x' \in K$ such that $Tx' = x'$. Since x' is also a minimizer of μ over E , it follows that for arbitrary $x \in E$ and for all $\xi \in (0, 1)$, $\Psi(x') \leq \Psi(x' + \xi(x - x'))$. Lemma 2.1 gives

$$(3.12) \quad \|z_n - x' - \xi(x - x')\|^q \leq \|z_n - x'\|^q - q\xi \left\langle x - x', j_q(z_n - x' - \xi(x - x')) \right\rangle.$$

Inequality (3.12) implies that

$$(3.13) \quad \mu_n \left\langle x - x', j_q(z_n - x' - \xi(x - x')) \right\rangle \leq 0.$$

Furthermore, since E is uniformly smooth, the generalized duality mapping j_q is norm-to-norm uniformly continuous on bounded subsets of E ; and hence

$$\lim_{\xi \rightarrow 0} \left(\left\langle x - x', j_q(z_n - x') \right\rangle - \left\langle x - x', j_q(z_n - x' - \xi(x - x')) \right\rangle \right) = 0.$$

So, for all $\epsilon > 0$, there exists $\delta_\epsilon > 0$ such that for all $\xi \in (0, \delta_\epsilon)$ and for all $n \in \mathbb{N}$, we have that

$$\left\langle x - x', j_q(z_n - x') \right\rangle - \left\langle x - x', j_q(z_n - x' - \xi(x - x')) \right\rangle < \epsilon.$$

This implies that for all $\xi \in (0, \delta_\epsilon)$,

$$\mu_n \left\langle x - x', j_q(z_n - x') \right\rangle \leq \epsilon + \mu_n \left\langle x - x', j_q(z_n - x' - \xi(x - x')) \right\rangle \leq \epsilon$$

and since $\epsilon > 0$ is arbitrary, we obtain that

$$\mu_n \left\langle x - x', j_q(z_n - x') \right\rangle \leq 0 \quad \forall x \in E.$$

Hence, we have in particular, that for the fixed $u \in E$ and $2x' - \lambda Ax' \in E$

$$(3.14) \quad \mu_n \left\langle u - x', j_q(z_n - x') \right\rangle \leq 0$$

and

$$(3.15) \quad \mu_n \left\langle (2x' - \lambda Ax') - x', j_q(z_n - x') \right\rangle = \mu_n \left\langle x' - \lambda Ax', j_q(z_n - x') \right\rangle \leq 0.$$

Moreover, since $\{t_n\}_{n \geq 1}$ is in $\left(0, \min\left\{1, \frac{1}{q\eta\lambda - d_q(\lambda L)^q}\right\}\right)$ and $x' \in F(T)$, we obtain from (3.10) that

$$\begin{aligned} \|z_n - x'\|^q &\leq \frac{1}{q\eta\lambda - d_q(\lambda L)^q} \left\langle u - \lambda Ax', j_q(z_n - x') \right\rangle \\ (3.16) \quad &= \frac{1}{\omega} \left(\left\langle u - x', j_q(z_n - x') \right\rangle + \left\langle x' - \lambda Ax', j_q(z_n - x') \right\rangle \right), \end{aligned}$$

where $\omega = q\eta\lambda - d_q(\lambda L)^q$. So, from (3.16) (using (3.14), (3.15) and the linearity of Banach limit) we obtain that

$$\begin{aligned} \mu_n \|z_n - x'\|^q &\leq \frac{1}{\omega} \mu_n \left\langle u - x', j_q(z_n - x') \right\rangle + \\ (3.17) \quad &+ \frac{1}{\omega} \mu_n \left\langle x' - \lambda Ax', j_q(z_n - x') \right\rangle \leq 0. \end{aligned}$$

So, $\mu_n \|z_n - x'\|^q = 0$ and this implies that there exists a subsequence $\{z_{n_i}\}_{i \geq 1}$ of $\{z_n\}_{n \geq 1}$ such that $z_{n_i} \rightarrow x'$ as $i \rightarrow \infty$.

We now show that x' is a solution of the variational inequality (3.7). From (3.3), we obtain that

$$z_{n_i} = t_{n_i}u + (I - t_{n_i}\lambda A)Tz_{n_i}.$$

This gives

$$(3.18) \quad \lambda Az_{n_i} - u = -\frac{1}{t_{n_i}}(z_{n_i} - Tz_{n_i}) + \lambda(Az_{n_i} - ATx_{n_i}).$$

Thus, for all $p \in F(T)$, we obtain from (3.18) that

$$\begin{aligned} \left\langle \lambda Az_{n_i} - u, j_q(z_{n_i} - p) \right\rangle &= -\frac{1}{t_{n_i}} \left\langle z_{n_i} - Tz_{n_i}, j_q(z_{n_i} - p) \right\rangle \\ (3.19) \quad &+ \lambda \left\langle Az_{n_i} - ATx_{n_i}, j_q(z_{n_i} - p) \right\rangle \end{aligned}$$

Since T is nonexpansive, we have that $(I - T)$ is accretive. Thus,

$$\left\langle z_{n_i} - Tz_{n_i}, j_q(z_{n_i} - p) \right\rangle = \left\langle (I - T)z_{n_i} - (I - T)p, j_q(z_{n_i} - p) \right\rangle \geq 0.$$

So, (3.19) gives

$$(3.20) \quad \left\langle \lambda Az_{n_i} - u, j_q(z_{n_i} - p) \right\rangle \leq \lambda \left\langle Az_{n_i} - ATx_{n_i}, j_q(z_{n_i} - p) \right\rangle$$

Using (3.20), we obtain (for some constant $M_2 > 0$) that

$$\begin{aligned} \left\langle \lambda Ax' - u, j_q(x' - p) \right\rangle &= \left\langle \lambda Az_{n_i} - u, j_q(z_{n_i} - p) \right\rangle \\ &\quad - \left\langle \lambda Az_{n_i} - \lambda Ax', j_q(z_{n_i} - p) \right\rangle \\ &\quad + \left\langle u - \lambda Ax', j_q(z_{n_i} - p) - j_q(x' - p) \right\rangle \\ &\leq M_2 (\|Az_{n_i} - ATz_{n_i}\| + \|\lambda Az_{n_i} - \lambda Ax'\| \\ (3.21) \quad &+ \|j_q(z_{n_i} - p) - j_q(x' - p)\|). \end{aligned}$$

Hence, since A, T are continuous, $z_{n_i} \rightarrow x' \in F(T)$ as $i \rightarrow \infty$ and the generalized duality mapping is norm-to-norm uniformly continuous on bounded subsets of E (and thus norm-to-norm continuous on bounded subsets of E), we obtain from (3.21) that (as $i \rightarrow \infty$)

$$\langle \lambda Ax' - u, j_q(x' - p) \rangle \leq 0 \quad \forall p \in F(T).$$

So, $x' \in F(T)$ is a solution (3.7).

Finally, we show that $\{z_n\}_{n \geq 1}$ converges to x' . Suppose that there is another subsequence $\{z_{n_l}\}_{l \geq 1}$ of $\{z_n\}_{n \geq 1}$ such that $z_{n_l} \rightarrow z' \in E$ as $l \rightarrow \infty$. Then by serial number (2) of Lemma 3.4, we have that $z' \in F(T)$. Similar argument (from (3.18) to (3.21) with z_{n_i} replaced by z_{n_l}) shows that

$$\langle \lambda Az' - u, j_q(z' - p) \rangle \leq 0 \quad \forall p \in F(T).$$

Uniqueness of solution of (3.7) shows that $x' = z'$. Hence, $z_n = z_{t_n} \rightarrow x'$ and $n \rightarrow \infty$. Consequently, we obtain that the path $\{z_t\}$ given by (3.3) converges strongly (as $t \rightarrow 0$) to unique $x' \in F(T)$ which solves the variational inequality (3.7). This completes the proof. $\square$

Remark 3.6. One may worry about how $\lim_{t \rightarrow 0} z_t = x'$. Actually, this follows from the fact that for any sequence $\{t_m\}_{m \geq 1}$ in $\left(0, \min\left\{1, \frac{1}{q\eta\lambda - d_q(L\lambda)^q}\right\}\right)$ such that $t_m \rightarrow 0$ as $m \rightarrow \infty$, the argument of the proof of Theorem 3.5 shows that $\{z_{t_m}\}_{m \geq 1} = \{z_m\}_{m \geq 1}$ converges to $x^* \in F(T)$ which solves the variational inequality (3.7) and since solution of (3.7) is unique, then $x^* = x'$. Thus, the path $\{z_t\}$ has a unique accumulation point as $t \rightarrow 0$.

The following corollaries follow from our discussion so far.

Corollary 3.7. *Let E be a real L_p or (ℓ_p) space. Let A and T be as in Lemma 3.3 and $\{z_t\}$ satisfy (3.3), then we obtain the same conclusion as in Theorem 3.5.*

Remark 3.8. Since every Hilbert space is a 2-uniformly smooth Banach space, it follows from Lemma 2.6 that if $E = H$ is a Hilbert space, then $d_q = d_2 = 1$. Furthermore, we recall that in a Hilbert space H , the duality mapping coincide with the identity mapping on H . Thus, we have the following corollary.

Corollary 3.9. *Let H be a real Hilbert space, $T : H \rightarrow H$ a nonexpansive mapping such that $F(T) \neq \emptyset$ and $A : H \rightarrow H$ be an L -Lipschitzian strongly accretive mapping with a constant $\eta > 0$. Let $u \in H$ be fixed. Suppose that $0 < \lambda < \frac{2\eta}{L^2}$ and $t \in \left(0, \min\left\{1, \frac{1}{2\eta\lambda - (L\lambda)^2}\right\}\right)$, then there exists a unique $z_t \in H$ satisfying (3.3). Moreover, $\{z_t\}$ converges strongly (as $t \rightarrow 0$) to a unique solution $x' \in F(T)$ of the variational inequality $\langle u - \lambda Ax', p - x' \rangle \leq 0 \quad \forall p \in F(T)$.*

Corollary 3.10. *Let E be a strictly convex q -uniformly smooth real Banach space, let $A : E \rightarrow E$ be an L -Lipschitzian strongly accretive mapping with a constant $\eta > 0$. Let $T_i : E \rightarrow E, i = 1, 2, \dots$ be a countable family of nonexpansive mappings such that $F := \bigcap_{i=1}^{\infty} F(T_i) \neq \emptyset$. Let $T := \sum_{i=1}^{\infty} \sigma_i T_i$, where $\{\sigma_i\}_{i \geq 1} \subset (0, 1)$ is such*

that $\sum_{i=1}^{\infty} \sigma_i = 1$. Let $u \in E$ be fixed. Suppose that the conditions of Lemma 3.3 are satisfied, then there exist a unique $z_t \in E$ satisfying $z_t = tu + (I - t\lambda A)Tz_t$. Moreover, $\{z_t\}$ converges strongly (as $t \rightarrow 0$) to a unique solution $x' \in F(T)$ of the variational inequality $\langle (u - \lambda Ax', j(p - x')) \rangle \leq 0 \forall p \in F$.

Proof. By Lemma 2.3, $T := \sum_{i=1}^{\infty} \sigma_i T_i$ is well defined, nonexpansive and $F(T) = F$. The rest follows as in the proof of Theorem 3.5. $\square$

Corollary 3.11. Let E be a strictly convex q -uniformly smooth real Banach space, let $A : E \rightarrow E$ be an L -Lipschitzian strongly accretive mapping with a constant $\eta > 0$. Let $T_i : E \rightarrow E, i = 1, 2, \dots, m$ be a finite family of nonexpansive mappings such that $F := \bigcap_{i=1}^m F(T_i) \neq \emptyset$. Let $T := \sum_{i=1}^m \sigma_i T_i$, where $\{\sigma_i\}_{i=1}^m \subset (0, 1)$ is such that $\sum_{i=1}^m \sigma_i = 1$. Let $u \in E$ be fixed. Suppose that the conditions of Lemma 3.3 are satisfied, then there exist a unique $z_t \in E$ satisfying $z_t = tu + (I - t\lambda A)Tz_t$. Moreover, $\{z_t\}$ converges strongly to a unique solution $x' \in F(T)$ of the variational inequality $\langle u - \lambda Ax', j(p - x') \rangle \leq 0 \forall p \in F$.

4. STRONG CONVERGENCE OF EXPLICIT ITERATION SCHEME FOR FINITE FAMILY OF NONEXPANSIVE MAPPINGS.

In the sequel, we shall assume that $0 < \lambda < \left(\frac{q\eta}{L^q d_q}\right)^{\frac{1}{q-1}}, \alpha_n \in \left(0, \min\left\{1, \frac{1}{q\eta\lambda - d_q(L\lambda)^q}\right\}\right)$

$\forall n \in \mathbb{N}, \lim_{n \rightarrow \infty} \alpha_n = 0, \sum_{n=1}^{\infty} \alpha_n = \infty$ and $\lim_{n \rightarrow \infty} \frac{\alpha_{n+r} - \alpha_n}{\alpha_{n+r}} = 0 \iff \lim_{n \rightarrow \infty} \frac{\alpha_n}{\alpha_{n+r}} = 1$.

Theorem 4.1. Let E be a q -uniformly smooth real Banach space which admits weakly sequentially continuous generalized duality mapping, let $A : E \rightarrow E$ be an L -Lipschitzian strongly accretive mapping with a constant $\eta > 0$. Let $T_i : E \rightarrow E, i = 1, 2, \dots, r$ be a finite family of nonexpansive mappings such that $\Omega := \bigcap_{i=1}^r F(T_i) \neq \emptyset$. Let $u \in E$ be fixed and $\{x_n\}_{n \geq 1}$ be a sequence in E generated iteratively by

$$(4.1) \quad x_0 \in E, x_{n+1} = \alpha_{n+1}u + (I - \alpha_{n+1}\lambda A)T_{n+1}x_n, n \geq 0,$$

where $T_n = T_{n \bmod r}$ and mod function takes values in $\{1, 2, \dots, r\}$. Suppose that $\Omega = F(T_r T_{r-1} \dots T_1) = F(T_1 T_{r-1} \dots T_2) = \dots = F(T_{r-1} \dots T_1 T_r)$, then, $\{x_n\}_{n \geq 0}$ converges strongly to a solution of the variational inequality

$$(4.2) \quad \langle u - \lambda Ax', j_q(p - x') \rangle \leq 0 \forall p \in \Omega.$$

Proof. Since the mapping $G = T_1 T_2 \dots T_r : E \rightarrow E$ is nonexpansive, then following the method of proof of Theorem 3.5, we have that for all $t \in \left(0, \min\left\{1, \frac{1}{q\eta\lambda - d_q(L\lambda)^q}\right\}\right)$, there exists unique $z_t \in E$ which satisfies $z_t = tu + (I - t\lambda A)Gz_t$, moreover, $\{z_t\}$

converges to a unique solution $x' \in F(G) = \Omega$ of (4.2). We now show that the explicit scheme (4.1) converges strongly to x' . We start by showing first that $\{x_n\}_{n \geq 0}$ is bounded. Now, let $p \in \Omega$ and set

$$r := \max \left\{ \|x_0 - p\|, \frac{q\|u - \lambda Ap\|}{q\eta\lambda - d_q(L\lambda)^q} \right\}.$$

We show by induction that

$$(4.3) \quad \|x_n - p\| \leq r \quad \forall n \geq 0.$$

Observe that for $n = 0$ (4.3) clearly holds. Assume for $n > 0$ that (4.3) is true. We show that (4.3) is also true for $n + 1$. Suppose for contradiction that this does not hold, then

$$\|x_{n+1} - p\| > r \geq \|x_n - p\|.$$

Thus, using (3.2), (4.1) and Lemma 2.1, we obtain that

$$\begin{aligned} \|x_{n+1} - p\|^q &= \|\alpha_{n+1}u + (I - \alpha_{n+1}\lambda A)T_{n+1}x_n - p\|^q \\ &= \|(I - \alpha_{n+1}\lambda A)T_{n+1}x_n \\ &\quad - (I - \alpha_{n+1}\lambda A)p + \alpha_{n+1}(u - \lambda Ap)\|^q \\ &\leq \|(I - \alpha_{n+1}\lambda A)T_{n+1}x_n - (I - \alpha_{n+1}\lambda A)p\|^q \\ &\quad + q\alpha_{n+1} \left\langle u - \lambda Ap, j_q(x_{n+1} - p) \right\rangle \\ &\leq \left(1 - \alpha_{n+1}(q\eta\lambda - d_q(L\lambda)^q)\right) \|x_n - p\|^q \\ &\quad + q\alpha_{n+1} \|u - \lambda Ap\| \cdot \|x_{n+1} - p\|^{q-1} \\ &< \left(1 - \alpha_{n+1}(q\eta\lambda - d_q(L\lambda)^q)\right) \|x_{n+1} - p\|^q \\ &\quad + q\alpha_{n+1} \|u - \lambda Ap\| \cdot \|x_{n+1} - p\|^{q-1}. \end{aligned} \quad (4.4)$$

Inequality (4.4) implies that

$$\|x_{n+1} - p\| < \frac{q\|u - \lambda Ap\|}{q\eta\lambda - d_q(L\lambda)^q},$$

a contradiction. Hence, the sequence $\{x_n\}_{n \geq 1}$ is bounded. Consequently, $\{T_{n+1}x_n\}_{n \geq 1}$ and $\{AT_{n+1}x_n\}_{n \geq 1}$ are also both bounded.

Furthermore, using (4.1),

$$\begin{aligned} x_{n+r} - x_n &= (\alpha_{n+r} - \alpha_n)u + (I - \alpha_{n+r}\lambda A)T_{n+r}x_{n+r-1} - (I - \alpha_{n+r}\lambda A)T_nx_{n-1} \\ &\quad - (\alpha_{n+r} - \alpha_n)\lambda AT_nx_{n-1}. \end{aligned} \quad (4.5)$$

Thus, using the fact that $T_{n+r} = T_n$, we obtain from (4.5) using Lemma 2.1 that for some constant $M_5 > 0$,

$$\begin{aligned} \|x_{n+r} - x_n\|^q &\leq \left(1 - \alpha_{n+1}(q\eta\lambda - d_q(L\lambda)^q)\right) \|x_{n+r-1} - x_{n-1}\|^q \\ &\quad + |\alpha_{n+r} - \alpha_n| M_5 \\ &= \left(1 - \alpha_{n+r}(q\eta\lambda - d_q(L\lambda)^q)\right) \|x_{n+r-1} - x_{n-1}\|^q \\ &\quad + M_5 \alpha_{n+r} \frac{|\alpha_{n+r} - \alpha_n|}{\alpha_{n+r}} \end{aligned} \quad (4.6)$$

So, using (4.6), we obtain from Lemma 2.2 that

$$(4.7) \quad \lim_{n \rightarrow \infty} \|x_{n+r} - x_n\| = 0$$

Furthermore, from the recursion formula (4.1) and for some constant $M_6 > 0$, we obtain

$$\|x_{n+1} - T_{n+1}x_n\| = \alpha_{n+1}\|u - \lambda AT_{n+1}x_n\| \leq \alpha_{n+1}M_6.$$

Thus,

$$\lim_{n \rightarrow \infty} \|x_{n+1} - T_{n+1}x_n\| = 0.$$

Similar argument shows that

$$(4.8) \quad x_{n+r} - T_{n+r}x_{n+r-1} \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Thus, we obtain using (4.8) and the fact that T_n is nonexpansive that

$$\begin{aligned} x_{n+r} - T_{n+r}x_{n+r-1} &\rightarrow 0 \text{ as } n \rightarrow \infty \\ T_{n+r}x_{n+r-1} - T_{n+r}T_{n+r-1}x_{n+r-2} &\rightarrow 0 \text{ as } n \rightarrow \infty \\ T_{n+r}T_{n+r-1}x_{n+r-2} - T_{n+r}T_{n+r-1}T_{n+r-2}x_{n+r-3} &\rightarrow 0 \text{ as } n \rightarrow \infty \\ &\vdots \\ T_{n+r}\dots T_{n+2}x_{n+1} - T_{n+r}\dots T_{n+2}T_{n+1}x_n &\rightarrow 0 \text{ as } n \rightarrow \infty. \end{aligned}$$

Adding up these yields

$$(4.9) \quad x_{n+r} - T_{n+r}\dots T_{n+2}T_{n+1}x_n \rightarrow 0 \text{ as } n \rightarrow \infty.$$

But,

$$(4.10) \quad \begin{aligned} \|x_n - T_{n+r}\dots T_{n+2}T_{n+1}x_n\| &\leq \|x_n - x_{n+r}\| \\ &\quad + \|x_{n+r} - T_{n+r}\dots T_{n+2}T_{n+1}x_n\| \end{aligned}$$

Thus, using (4.7), (4.9) and (4.10), we get that

$$(4.11) \quad \lim_{n \rightarrow \infty} \|x_n - T_{n+r}\dots T_{n+2}T_{n+1}x_n\| = 0.$$

Next, we show that

$$\limsup_{n \rightarrow \infty} \langle u - \lambda Ax', j_q(x_n - x') \rangle \leq 0.$$

Let $\{x_{n_k}\}_{k \geq 1}$ be a subsequence of $\{x_n\}_{n \geq 1}$ such that

$$\limsup_{n \rightarrow \infty} \langle u - \lambda Ax', j_q(x_n - x') \rangle = \lim_{k \rightarrow \infty} \langle u - \lambda Ax', j_q(x_{n_k} - x') \rangle.$$

Since $\{x_n\}_{n \geq 1}$ is bounded and since E is a reflexive real Banach space, there exists a subsequence $\{x_{n_{k_m}}\}_{m \geq 1}$ of $\{x_{n_k}\}_{k \geq 1}$ such that $\{x_{n_{k_m}}\}_{m \geq 1}$ converges weakly to some $p^* \in E$. Without loss of generality, we may assume that n_{k_m} is such that $T_{n_{k_m}} = T_i$ for some $i \in \{1, 2, \dots, r\}$, for all $m \geq 1$. It therefore follows from (4.11) that

$$\lim_{m \rightarrow \infty} \|x_{n_{k_m}} - T_{i+r}\dots T_{i+2}T_{i+1}x_{n_{k_m}}\| = 0.$$

So, by Lemma 2.7, $p^* \in F(T_{i+r} \dots T_{i+2} T_{i+1}) = \Omega$. Thus, since E has weakly sequential continuous generalized duality mapping, we have that

$$\begin{aligned}
 \limsup_{n \rightarrow \infty} \langle u - \lambda A x', j_q(x_n - x') \rangle &= \lim_{k \rightarrow \infty} \langle u - \lambda A x', j_q(x_{n_k} - x') \rangle \\
 &= \lim_{m \rightarrow \infty} \langle u - \lambda A x', j_q(x_{n_{k_m}} - x') \rangle \\
 &= \langle u - \lambda A x', j_q(p^* - x') \rangle \leq 0
 \end{aligned}
 \tag{4.12}$$

Thus, setting

$$\theta_n = \max \left\{ 0, \langle u - \lambda A x', j_q(x_n - x') \rangle \right\},$$

then it is easy to see that $\lim_{n \rightarrow \infty} \theta_n = 0$. Furthermore, we obtain from the recursion formula (4.1) using Lemma 2.1 that

$$\begin{aligned}
 \|x_{n+1} - x'\|^q &\leq \left(1 - \alpha_{n+1}(q\eta\lambda - d_q(\lambda L)^q) \right) \|x_n - x'\|^q \\
 &\quad + q\alpha_{n+1} \langle u - \lambda A x', j_q(x_{n+1} - x') \rangle \\
 &\leq \left(1 - \alpha_{n+1}(q\eta\lambda - d_q(\lambda L)^q) \right) \|x_n - x'\|^q + \delta_n \quad \forall n \geq 0,
 \end{aligned}$$

where $\delta_n = \alpha_{n+1}q(q\eta\lambda - d_q(\lambda L)^q) \left[\frac{\theta_{n+1}}{q\eta\lambda - d_q(\lambda L)^q} \right]$, which is clear $o(\alpha_{n+1})$. Hence, by Lemma 2.2, $\{x_n\}_{n \geq 1}$ converges strongly to $x' \in \Omega$ which is a unique solution of (4.2). This completes the Proof. $\square$

Remark 4.2. We note that in the corresponding result of Xu [48], the assumption

$$\Omega = F(T_r T_{r-1} \dots T_1) = F(T_1 T_{r-1} \dots T_2) = \dots = F(T_{r-1} \dots T_1 T_r)$$

was made. We now consider a situation where this condition is dispensed with.

Example 4.3. Let E be a strictly convex q -uniformly smooth real Banach space which admits weakly sequentially continuous generalized duality mapping, let $A : E \rightarrow E$ be an L -Lipschitzian strongly accretive mapping with a constant $\eta > 0$. Let $T_i : E \rightarrow E, i = 1, 2, \dots, r$ be a finite family of nonexpansive mappings such that $\Omega := \bigcap_{i=1}^r F(T_i) \neq \emptyset$ and $S_i = (1 - \omega_i)I + \omega_i T_i, i = 1, 2, \dots, r$. Let $u \in E$ be fixed and let $\{x_n\}_{n \geq 1}$ be a sequence in E generated iteratively by

$$(4.13) \quad x_0 \in E, x_{n+1} = \alpha_{n+1}u + (I - \alpha_{n+1}\lambda A)S_{n+1}x_n, \quad n \geq 0,$$

where $S_n = S_{n \bmod r}$ and mod function takes values in $\{1, 2, \dots, r\}$, then $\{x_n\}_{n \geq 0}$ converges strongly to a solution of the variational inequality (4.2)

Proof. By Lemma 2.4, $\bigcap_{i=1}^r F(S_i) = \bigcap_{i=1}^r F(T_i)$ and $\bigcap_{i=1}^r F(S_i) = F(S_r S_{r-1} \dots S_1) = F(S_1 S_{r-1} \dots S_2) = \dots = F(S_{r-1} \dots S_1 S_r)$. The rest follows as in the proof of Theorem 4.1. $\square$

Corollary 4.4. Let E be a q -uniformly smooth real Banach space which admits weakly sequentially continuous generalized duality mapping, let $A : E \rightarrow E$ be an L -Lipschitzian strongly accretive mapping with a constant $\eta > 0$. Let $T : E \rightarrow E$,

be a nonexpansive mapping such that $F(T) \neq \emptyset$. Let $u \in E$ be fixed and let $\{x_n\}_{n \geq 1}$ be a sequence in E generated iteratively by

$$(4.14) \quad x_0 \in E, x_{n+1} = \alpha_{n+1}u + (I - \alpha_{n+1}\lambda A)Tx_n, \quad n \geq 0,$$

then $\{x_n\}_{n \geq 0}$ converges strongly to a unique $x' \in F(T)$ which is a solution of the variational inequality (3.7)

Proof. Follows as in the proof of Theorem 4.1 using Theorem 3.5. $\square$

Remark 4.5. We remark that corollaries synonymous to Corollaries 3.7, 3.9, 3.10 and 3.11 are obtainable in this section. But we must note, however, that though Theorem 4.1, Corollary 4.3 and corollary 4.4 hold in the sequence space ℓ^q , they do not hold in $L^q(\mathbb{R})$ for $1 < q < +\infty$, $q \neq 2$ since $L^q(\mathbb{R})$, $q \neq 2$ do not possess weakly sequentially continuous duality mapping.

Remark 4.6. The addition of bounded error terms to our recursion formulas leads to no further generalization. It is easy to see that if we replace u by λu in both implicit and explicit iteration scheme studied in this paper, we obtain that our iteration schemes converge strongly to a unique solution $x' \in \Omega$ of the variational inequality $\langle u - Ax', j_q(p - x') \rangle \leq 0 \quad \forall p \in \Omega$, where Ω is the set of common fixed points of finite family of nonexpansive mappings.

5. APPLICATIONS

CONVERGENCE THEOREM FOR FAMILIES OF STRICTLY PSEUDOCONTRACTIVE MAPPINGS.

Let E be a normed space. A mapping $T : E \rightarrow E$ is called *k-strictly pseudocontractive* if and only if there exists a real constant $k > 0$ such that for all $x, y \in D(T)$ there exists $j(x - y) \in J(x - y)$ such that

$$(5.1) \quad \langle Tx - Ty, j(x - y) \rangle \leq \|x - y\|^2 - k\|x - y - (Tx - Ty)\|^2.$$

Without loss of generality we may assume that $k \in (0, 1)$. If I denotes the identity operator, then (5.1) can be re-written as

$$(5.2) \quad \langle (I - T)x - (I - T)y, j(x - y) \rangle \geq k\|(I - T)x - (I - T)y\|^2.$$

In Hilbert spaces, (5.1) (or equivalently (5.2)) is equivalent to the inequality

$$\|Tx - Ty\|^2 \leq \|x - y\|^2 + \beta\|(I - T)x - (I - T)y\|^2, \quad \text{where } \beta = (1 - k) < 1.$$

It was shown in [33] that if T is *k-strictly pseudocontractive*, then the following inequality holds

$$(5.3) \quad \langle (I - T)x - (I - T)y, j_q(x - y) \rangle \geq k^{q-1}\|(I - T)x - (I - T)y\|^q.$$

Thus, if E is a q -uniformly smooth real Banach space; and $T : E \rightarrow E$ is a *k-strictly pseudocontractive* mapping, then for the map $T_a := (1 - a)I + aT : E \rightarrow E$ (where I is the identity map of E and $a > 0$), we obtain by Lemma 2.5 using (5.3) that:

$$\begin{aligned} \|T_ax - T_ay\|^q &= \|x - y - a((I - T)x - (I - T)y)\|^q \\ &\leq \|x - y\|^q - qa\langle (I - T)x - (I - T)y, j_q(x - y) \rangle \\ &\quad + d_q a^q \|(I - T)x - (I - T)y\|^q \\ &\leq \|x - y\|^q - a(k^{q-1}q - d_q a^{q-1})\|Ax - Ay\|^q, \end{aligned}$$

where $A = (I - T)$. If a is such that $0 < a < \left(\frac{qk^{q-1}}{d_q}\right)^{\frac{1}{q-1}}$, we have that the mapping T_a is nonexpansive. It is also easy to see that the fixed point set of T_a and that of T coincide.

Thus, we have the following theorem

Theorem 5.1. *Let E be a q -uniformly smooth real Banach space which admits weakly sequential continuous generalized duality mapping, let $A : E \rightarrow E$ be an L -Lipschitzian strongly accretive mapping with a constant $\eta > 0$. Let $T_i : E \rightarrow E, i = 1, 2, \dots, r$ be a finite family of k -strictly pseudocontractive mappings such that $\Omega^* := \bigcap_{i=1}^r F(T_i) \neq \emptyset$. Let $\{a_i\}_{i=1}^r$ be such that $0 < a_i < \left(\frac{qk^{q-1}}{d_q}\right)^{\frac{1}{q-1}}, i = 1, 2, \dots, r$ and define $T_{a_i} = (1 - a_i)I + a_i T_i$. Let $u \in E$ be fixed and $\{x_n\}_{n \geq 1}$ be a sequence in E generated iteratively by*

$$(5.4) \quad x_0 \in E, x_{n+1} = \alpha_{n+1}u + (I - \alpha_{n+1}\lambda A)T_{a_{n+1}}x_n, \quad n \geq 0,$$

where $T_{a_n} = T_{a_{n \bmod r}}$ and mod function takes values in $\{1, 2, \dots, r\}$. Suppose that $\Omega^* = F(T_{a_r}T_{a_{r-1}}\dots T_{a_1}) = F(T_{a_1}T_{a_{r-1}}\dots T_{a_2}) = \dots = F(T_{a_{r-1}}\dots T_{a_1}T_{a_r})$, then, $\{x_n\}_{n \geq 0}$ converges strongly to a solution of the variational inequality

$$(5.5) \quad \langle u - \lambda Ax', j_q(p - x') \rangle \leq 0 \quad \forall p \in \Omega^*.$$

Corollary 5.2. *Let E be a strictly convex q -uniformly smooth real Banach space which admits weakly sequentially continuous generalized duality mapping, let $A : E \rightarrow E$ be an L -Lipschitzian strongly accretive mapping with a constant $\eta > 0$. Let $T_i : E \rightarrow E, i = 1, 2, \dots, r$ be a finite family of k -strictly pseudocontractive mappings such that $\Omega := \bigcap_{i=1}^r F(T_i) \neq \emptyset$. Let $\{a_i\}_{i=1}^r$ be such that $0 < a_i < \left(\frac{qk^{q-1}}{d_q}\right)^{\frac{1}{q-1}}, i = 1, 2, \dots, r$ and define $T_{a_i} = (1 - a_i)I + a_i T_i$ and $S_{a_i} = (1 - \omega_i)I + \omega_i T_{a_i}, i = 1, 2, \dots, r$. Let $u \in E$ be fixed and let $\{x_n\}_{n \geq 1}$ be a sequence in E generated iteratively by*

$$(5.6) \quad x_0 \in E, x_{n+1} = \alpha_{n+1}u + (I - \alpha_{n+1}\lambda A)S_{a_{n+1}}x_n, \quad n \geq 0,$$

where $S_{a_n} = S_{n \bmod r}$ and mod function takes values in $\{1, 2, \dots, r\}$. Then, $\{x_n\}_{n \geq 0}$ converges strongly to a solution of the variational inequality (5.5).

Corollary 5.3. *Let E be a q -uniformly smooth real Banach space which admits weakly sequentially continuous generalized duality mapping, let $A : E \rightarrow E$ be an L -Lipschitzian strongly accretive mapping with a constant $\eta > 0$. Let $T : E \rightarrow E$, be a k -strictly pseudocontractive mapping such that $F(T) \neq \emptyset$. Let be such that $0 < a < \left(\frac{qk^{q-1}}{d_q}\right)^{\frac{1}{q-1}}$ and define $T_a = (1 - a)I + aT$. Let $u \in E$ be fixed and let $\{x_n\}_{n \geq 1}$ be a sequence in E generated iteratively by*

$$(5.7) \quad x_0 \in E, x_{n+1} = \alpha_{n+1}u + (I - \alpha_{n+1}\lambda A)T_ax_n, \quad n \geq 0,$$

then, $\{x_n\}_{n \geq 0}$ converges strongly to a unique $x' \in F(T)$ which is a solution of the variational inequality (3.7)

Remark 5.4. Prototype for our iteration parameter $\{\alpha_n\}_{n \geq 1}$ (see e.g. [48]) is given by

$$\alpha_n = \begin{cases} \frac{1}{\sqrt{n}} & \text{if } n \text{ is odd,} \\ \frac{1}{\sqrt{n-1}} & \text{if } n \text{ is even.} \end{cases}$$

If we assume that r is odd, since the case r being even is similar, it is not difficult to see that

$$\frac{\alpha_n}{\alpha_{n+r}} = \begin{cases} \frac{\sqrt{n+r}-1}{\sqrt{n}} & \text{if } n \text{ is odd,} \\ \frac{\sqrt{n+r}}{\sqrt{n-1}} & \text{if } n \text{ is even.} \end{cases}$$

Remark 5.5. It is easy to see that Corollary 4.3 is an obvious improvement on the corresponding results of Xu [48] in the sense that the condition

$$\Omega = F(T_r T_{r-1} \dots T_1) = F(T_1 T_{r-1} \dots T_2) = \dots = F(T_{r-1} \dots T_1 T_r)$$

which was imposed by Xu is dispensed with.

Remark 5.6. When $E = H$, a Hilbert space, $\lambda = 1$ and A is a bounded linear strongly positive operator, our iteration process (4.1) reduces to the iteration scheme studied by Xu [48]. If the fixed vector $u \in E$ is identically equal to the zero vector of H , A a strongly monotone operator and we consider a single nonexpansive mapping, (4.1) reduces to the scheme studied by Yamada [50]. We recall that the results of Xu [48] and Yamada [50] remain in Hilbert spaces. Our theorems, therefore, extend, generalize, improve and unify the corresponding results of these authors and that of a host of other authors in the more general setting of q -uniformly real Banach spaces and for the more general class of strongly accretive mappings. Our corollaries, applications and method of proof are of independent interest.

REFERENCES

- [1] A. Aleyner and S. Reich, An explicit construction of sunny nonexpansive retractions in Banach spaces, *Hindawi Publishing Corporation-Fixed Point Theory and Applications*, 2005:3 (2005), 295-305, DOI:10.1155/FPTA.2005.295.
- [2] H. H. Bauschke, The Approximation of fixed points of compositions of nonexpansive mappings in Hilbert spaces, *J. Math. Anal. Appl.* 202 (1996) 150-159.
- [3] T. D. Benavides, G. L. Acedo and H. K. Xu, Construction of sunny nonexpansive retractions in Banach spaces, *Bull. Austral. Math. Soc.* vol. 66 (2002), 9-16.
- [4] V. Berinde, Iterative approximation of fixed points, *Editura Efemeride Baia Mare*, 2002.
- [5] F. E. Browder, Convergence theorems for sequences of nonlinear operators in Banach spaces, *Math. Zeitschr.* 100 (1967) 201-225.
- [6] F. E. Browder, Fixed point theorems for nonlinear semicontractive mappings in Banach spaces, *Archive for Rat. Mech. and Anal.* 21 (1966) 259-269.
- [7] F. E. Browder and D. G. De Figueirado, J -monotone nonlinear operators in Banach spaces, *Konk. Nederl. Akadi. Wetesch.* 69 (1966) 412-420.
- [8] R. E. Bruck; Jr. Nonexpansive projections on subsets of Banach space, *Pacific J. Math.* **47** (1973), 341-355.
- [9] R. E. Bruck, JR., Nonexpansive Retracts of Banach spaces, *Bull. Amer. Math. Soc.*, 76 (1970), 384-386.
- [10] R. E. Bruck, JR., Nonexpansive projections on subsets of Banach spaces, *Pacific Journal of Mathematics*, vol. 47, no.2 (1973), 341-355.
- [11] R. E. Bruck, Jr., Properties of Fixed point sets of nonexpansive mappings in Banach spaces, *Trans. Amer. Math. Soc.* **179** (1973), 251-262.

- [12] C. Byrne, A unified treatment of some iterative algorithms in signal processing and image construction, *Inverse problems* **20** (2004) 103-120.
- [13] C. E. Chidume, Geometric properties of Banach spaces and nonlinear iterations, *Springer Verlag Series: Lecture Notes in Mathematics*, Vol. 1965 (2009), ISBN 978-1-84882-189-7.
- [14] C. E. Chidume and E. U. Ofoedu, A new iteration process for finite families of generalized Lipschitz pseudo-contractive and generalized Lipschitz accretive mappings, *Nonlinear Analysis; TMA.*, In press, accepted manuscript, available online 27 June 2007.
- [15] C. E. Chidume and E. U. Ofoedu, Approximation of common fixed points for finite families of total asymptotically nonexpansive mappings, *J. Math. Anal. Appl.*, **333**(1) (2007), 128-141.
- [16] F. Deutch, I. Yamada, Minimizing certain convex functions over the intersection of the fixed point sets of nonexpansive mappings, *Numerical Functional Analysis and Optimization* **19** (1998) 33-56.
- [17] J. P. Gossez and E. Lami Dozo, Some geometric properties related to fixed point theory for nonexpansive mappings, *Pacific J. Math.* **40** (1972), 565-573.
- [18] B. Halpern, Fixed points of nonexpanding maps, *Bulletin of the American Mathematical Society*, **73** (1967), 957-961.
- [19] H. Iiduka and W. Takahashi, Strong convergence theorems for nonexpansive mappings and inverse-strongly monotone mappings, *Nonlinear Anal.* **61** (2005), 341-350.
- [20] S. Ishikawa, Fixed point by a new iteration method, *Proc. Amer. Math. Soc.* **44** (1974), 147-150.
- [21] T. C. Lim, H. K. Xu and Z. B. Xu, Some L_P inequalities and their applications to fixed point theory and approximation theory, in *progress in Approximation theory* (P. Nevai and A. Pinkus, Eds.), pp. 609-624, Academic press, 1991.
- [22] M. Lin and R. Sine, Contractive projections on the fixed point set of L_∞ contractions, *Colloquium Mathematicum*, vol. LXII (1991) FASC1.
- [23] P. L. Lions, Approximation de points fixes de contractions, *Comptes rendus de l'academie des sciences, serie I-mathematique*, **284** (1997) 1357-1359.
- [24] Y. Liu, A general iterative method for equilibrium problems and strict pseudo-contractions in Hilbert spaces, *Nonlinear Analysis, TMA* **71** (2009) 4852-4861.
- [25] W. R. Mann, Mean value methods in iteration, *Proc. Amer. Math. Soc.* **4** (1953) 506-510.
- [26] G. Marino, H. K. Xu, A general iterative method for nonexpansive mappings in Hilbert spaces, *J. Math. anal. Appl.* **318** (2006) 43-52.
- [27] R. E. Megginson, An introduction to Banach space theory, *Springer-Verlag*, New York, Inc., 1998.
- [28] A. Moudafi, Viscosity approximation methods for fixed-points problems, *J. Math. Anal. Appl.* **241** (2000) 46-55.
- [29] E. U. Ofoedu, Iterative approximation of a common zero of a countably infinite family of m -accretive operators in Banach spaces, *Hindawi-Fixed point theory and applications*, doi:10.1155/2008/325792.
- [30] E. U. Ofoedu and Y. Shehu, Iterative construction of a common fixed point of finite families of nonlinear mappings, in press, to appear in Vol. 16 No 1 of *Journal of Applied Analysis* (2010).
- [31] E. U. Ofoedu, Y. Shehu and J. N. Ezeora, Solution by iteration of nonlinear variational inequalities involving finite family of asymptotically nonexpansive mappings and monotone mappings, *PanAmer. Math. J.* **18** (2008) (4), 61-75.
- [32] Z. Opial, Weak convergence of the sequence of successive approximations for nonexpansive mappings, *Bulletin of the Australian Mathematical Society* **73** (1967), 591-597.
- [33] M. O. Osilike and A. Udomene, Demiclosedness principle and convergence theorems for strictly pseudocontractive mappings of Browder-Petryshyn type, *J. Math. Anal. Appl.* **256** (2001), 431-445.
- [34] C. I. Polilchuk and R. J. Mammone, Image recovery by convex projections using a least-square constraint, *J. Opt. Soc. Amer.* **A7** (1990) 517-521.
- [35] S. Reich, Weak convergence theorems for nonexpansive mappings in Banach spaces, *J. Math. Anal. Appl.* **67** (1979), 274-276.
- [36] S. Reich, Extension problems for accretive sets in Banach spaces, *J. Functional Analysis* **26** (1977), 378-395.
- [37] S. Reich, Strong convergence theorems for resolvents of accretive operators in Banach spaces, *J. Math. Anal. Appl.*, **75** (1980), 287-292.

- [38] B. E. Rhoades, Fixed point iterations for certain nonlinear mappings, *J. Math. Anal. Appl.* **183** (1994) 118-120.
- [39] R. T. Rockafellar, Monotone operators and proximal point algorithm, SIAM, *J. Control Optim.* **14** (1976), 877-898.
- [40] J. Schu, Iterative construction of fixed points of asymptotically nonexpansive mappings, *J. Math. Anal. Appl.* **158** (1991), 407-413.
- [41] J. Schu, Weak and strong convergence of fixed points of asymptotically nonexpansive mappings, *Bull. Austral. Math. Soc.* **43** (1991), 153-159.
- [42] S. Shioji, W. Takahashi; Strong convergence of approximated sequences for nonexpansive mappings in Banach spaces, *Proc. Amer. Math. Soc.* **125** (1997), 3641-3645.
- [43] M. Tian, A general iterative algorithm for nonexpansive mappings in Hilbert spaces, *Nonlinear Analysis (2010)*, doi: 10.1016/j.na.2010.03.058.
- [44] R. Wittmann, Approximation of fixed points of nonexpansive mappings, *Archiv der mathematik*, 58 (1992) 486-491.
- [45] D. Wang, Y. Song and X. Ma, Iterative methods for finite family of nonexpansive mappings, *Int. J. Contemp. Math. Sciences*, vol. 3 (2008) no. 9, 1409-1417.
- [46] H. K. Xu, Iterative algorithms for nonlinear operators, *J. London Math. Soc.* **66** (2002) 240-256.
- [47] H. K. Xu, Inequality in Banach spaces with applications, *Nonlinear Anal.* **16** (1991), 1127-1138, *London Math. Soc.*, **66** (2) (2002), 240-256.
- [48] H. K. Xu, An iterative approach to quadratic optimization, *Journal of Optimization Theory and Applications*, **116** (2003) no.3, 659-678.
- [49] H. K. Xu, Viscosity approximation methods for nonexpansive mappings, *J. Math. Anal. Appl.* **298** (2004) 279-291.
- [50] I. Yamada, The hybrid steepest decent for the variational inequality problems over the intersection of fixed points sets of nonexpansive mapping: in **D. Butnariu, Y. Censor, S. Reich (Eds)**, *Inherently parallel algorithms in feasibility and optimization and their application*, Elsevier, New York, (2001) 473-504.
- [51] D. Youla, On deterministic convergence of iterations of related projection operators, *J. Vis. Commun. Image Represent* **1** (1990), 12-20.
- [52] W. Takahashi, *Nonlinear Functional Analysis*, Yokohama publishers, Yokohama, 2000.
- [53] H. Zhou, L. Wei, Y. J. Cho; Strong convergence theorems on an iterative method for family of finite nonexpansive mappings in reflexive Banach spaces, *Appl. Math. Comput.* **173** (2006) 196-212.

¹DEPARTMENT OF MATHEMATICS, NNAMDI AZIKIWE UNIVERSITY, P. M. B. 5025, AWKA, ANAMBRA STATE, NIGERIA.

E-mail address: euofoedu@yahoo.com